


EFFECTS OF REDD INUNDATION RESULTING FROM RESERVOIR FLUCTUATION ON BULL TROUT (*SALVELINUS CONFLUENTUS*) EGG SURVIVAL AND EMERGENCE IN THE CEDAR RIVER WATERSHED, WASHINGTON

Dudley Reiser, Eric Jeanes, Stuart Beck,
Catherine Morello
R2 Resource Consultants
and
Dwayne Paige and Heidy Barnett
Seattle Public Utilities



Background

- ◆ Chester Morse Lake – major water supply for City of Seattle – also some limited hydropower generation
- ◆ Major tributaries – Cedar and Rex rivers




more background


- ◆ Cedar watershed supports Adfluvial populations of bull trout (“threatened” under ESA)
- ◆ Estimated population of bull trout: 3100 adults
- ◆ Pygmy whitefish (native) and rainbow trout (introduced) also present




and even more background

- ◆ Reservoir levels fluctuate seasonally
- ◆ Low levels create riverine habitats in lower reaches of rivers – conducive for spawning
- ◆ Higher levels – progressive upstream shift from riverine to lacustrine habitats – potential redd inundation








Bull trout spawning on rising limb of hydrograph



Bull trout spawning when reservoir levels low

Egg incubation success depends on:

- ◆ Suitable depths and velocities
- ◆ Intragravel velocities & downwelling
- ◆ Clean gravels = low fine sediments (<0.84 mm)
- ◆ DO





Lower redds 10-15 ft deep



City of Seattle Multi-species Habitat Conservation Plan -2000

- ◆ Assessment of potential bull trout “take”
required in HCP

ENDANGERED SPECIES ACT
- SECTION 7
BIOLOGICAL OPINION
UNLISTED SPECIES
ANALYSIS, AND SECTION 10
FINDINGS

for proposed issuance of a
Section 10 Incidental Take
Permit to the City of Seattle,
Seattle Public Utility, for the
Cedar River Watershed
Habitat Conservation Plan



And so....Study

- ◆ Effects of reservoir fluctuations on bull trout spawning and egg survival:
SPECIFICALLY – effects of redd inundation
Ho – Higher egg mortality related to longer periods of redd inundation
- ◆ Estimate potential “take” (if possible)
- ◆ Define causal mechanisms of take
- ◆ Integrate results into “risk-averse” operational plan

In-situ egg survival study

- ♦ Remote spawning of bull trout
- ♦ Artificial redds constructed within zone of reservoir influence
- ♦ Monitor redds and “controls” for survival and emergence
- ♦ Collect ancillary data
 - ♦ Sediment
 - ♦ Water temperature
 - ♦ DO
- ♦ Comparisons of Survival to emergence (STE) vs Days of inundation



Capture





Spawn – Fertilize – Water Harden



Fish stats

- ◆ Cedar River
 - ◆ 5 females spawned (400-528 mm FL)
 - ◆ 2 males (460-520 mm FL)
- ◆ Rex River
 - ◆ 5 females spawned (453-563 mm FL)
 - ◆ 1 male (546mm FL)
- ◆ Weights – range 650 – 1500 gm



Egg Planting



Egg chambers = Whitlock – Vibert boxes



- Filled with clean gravel
- Duct tape on bottom
- 50 embryos per box




- 4 boxes per redd
- Colored string attached for relocation

Cedar River Inundation Sites 2006

N=23

Approximate upper
extent of inundation

0 50 100 200 300 400 Meters




Rex River Inundation Sites 2006

N=17

Approximate upper
extent of inundation

0 55 110 220 330 440 Meters






CONTROLS



And then Mr. Murphy showed up – RAIN,RAIN,RAIN...



Effects

- ◆ Scour and dislodgement
- ◆ Deposition and covering
- ◆ Channel re-alignment



TESTS COMPROMISED

Cedar River Inundation Sites 2006

N=23


Not Found/Recovered

4% survival

Approximate upper
extent of inundation

76% survival

0 50 100 200 300 400 Meters




Rex River Inundation Sites 2006

N=17

Not Found/Recovered

Approximate upper
extent of inundation

0 55 110 220 330 440 Meters



PRELIMINARY POSTULATES:

- Observations
- Some physical data
- Some biological data






OBSERVATIONS

**Surficial sediment
deposition noticeable
at reservoir – river interface**



PHYSICAL DATA



BIOLOGICAL DATA


(yes it is limited): **spatial difference in survival**

- ◆ Riverine - Low sediment levels = high egg survival


- ◆ Lacustrine - High sediment levels = low egg survival



Operations - Static




Operations - Dynamic






**Risk of DEPOSITION
& ENTOMBMENT**

**Risk of low DO
& App Vel**

**Combinations of River Flow and
Reservoir Levels are complicated**



ANOTHER YEAR OF STUDY



QUESTIONS

